



## **Alteration of nephrocystins and IFT-A proteins causes similar ciliary phenotypes leading to Nephronophthisis**

Sophie Saunier, Albane A. Bizet, Flora Silbermann, Emilie Filhol, Thierry Blisnick, Anthony Henneveu, Emilie Montenont, Isabelle Perrault, Christine Boyle-Feysot, Jean-Michel Rozet, et al.

### **► To cite this version:**

Sophie Saunier, Albane A. Bizet, Flora Silbermann, Emilie Filhol, Thierry Blisnick, et al.. Alteration of nephrocystins and IFT-A proteins causes similar ciliary phenotypes leading to Nephronophthisis. First International Cilia in Development and Disease Scientific Conference, May 2012, London, United Kingdom. pp.P99, 10.1186/2046-2530-1-S1-P99 . inserm-00752970

**HAL Id: inserm-00752970**

**<https://www.hal.inserm.fr/inserm-00752970>**

Submitted on 16 Nov 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

# Alteration of nephrocystins and IFT-A proteins causes similar ciliary phenotypes leading to Nephronophthisis

S Saunier<sup>1\*</sup>, AA Bizet<sup>1</sup>, F Silbermann<sup>1</sup>, E Filhol<sup>1</sup>, T Blisnick<sup>2</sup>, A Henneveu<sup>3</sup>, E Montenont<sup>1</sup>, I Perrault<sup>4</sup>, C Boyle-Feysot<sup>5</sup>, J-M Rozet<sup>4</sup>, P Bastin<sup>2</sup>, HH Arts<sup>6</sup>, C Antignac<sup>1</sup>, AR Benmerah<sup>3</sup>

From First International Cilia in Development and Disease Scientific Conference (2012)  
London, UK. 16-18 May 2012

Nephronophthisis (NPH) is a kidney ciliopathy often associated with extra-renal defects and for which 12 genes (*NPHP1-12*) have been identified. *NPHP1* and *NPHP4* control the ciliary access at the transition zone and the velocity of some intraflagellar transport (IFT)/BBS proteins in *C.elegans*. Recently, in a collaborative effort, we have identified, in families with isolated NPH, mutations in *TTC21B* as well as in *WDR19*, which encode the retrograde IFT-A proteins IFT139 and IFT144, respectively. By ciliome sequencing of 1600 candidate genes from 14 NPH patients followed by Sanger sequencing of a cohort of 52 patients, we have found respectively 8 and 7 patients carrying pathogenic missense mutations in genes coding IFT-A proteins, including *WDR35*, *TTC21B* and *IFT140*, which could partially affect their function. Together, these results indicate that IFT-A are involved in nephronophthisis. Moreover, alteration of cilia length was observed in patient kidney, *Nphp4*<sup>-/-</sup> mice kidney tubules and *NPHP1* or *NPHP4* knockdown IMCD3 cell lines. In these cells, primary cilia present swellings at the distal region accompanied by an accumulation of IFT-B at the base and the tip, similar to what was observed in IFT-A mutants, suggesting a possible alteration of retrograde transport. Additionally, ARL13B, a small GTPase required for proper cilium shape and IFT stability, is absent along the axoneme of *NPHP4*-KD-IMCD cells. By controlling the entry of ciliary components at the transition zone, *NPHP1* and *NPHP4* may modulate IFT-A cargos thus participating in the same pathway (i.e. Wnt/PCP), alteration of which would lead to renal lesions observed in nephronophthisis.

## Author details

<sup>1</sup>Inserm U983, France. <sup>2</sup>Pasteur Institute, CNRS UMR2581, France.

<sup>3</sup>Inserm U1016, Cochin Institute, France. <sup>4</sup>Inserm U781, France.

<sup>5</sup>Fondation Imagine, Paris, France. <sup>6</sup>Radboud University Nijmegen Medical Center, the Netherlands.

Published: 16 November 2012

doi:10.1186/2046-2530-1-S1-P99

**Cite this article as:** Saunier et al.: Alteration of nephrocystins and IFT-A proteins causes similar ciliary phenotypes leading to Nephronophthisis. *Cilia* 2012 **1**(Suppl 1):P99.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)



\* Correspondence: [sophie.saunier@inserm.fr](mailto:sophie.saunier@inserm.fr)

<sup>1</sup>Inserm U983, France

Full list of author information is available at the end of the article